

External



Ahmad Dwi Setyawan <smujo.id@gmail.com>

Mon, Apr 25, 2:15 PM

to me

Ni Putu Pandawani:

Thank you for submitting the manuscript, "High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection" to Biodiversitas Journal of Biological Diversity. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Submission URL: <https://smujo.id/biodiv/authorDashboard/submission/11071>

Username: pandawani

11097 / **Pandawani** / High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection

Upload FileLibrary

WorkflowPublication

- [Submission](#)
-

- [Review](#)
-

- **Copyediting**

- **Production**

Submission Files

- **Search**

Name	Date	Component
Settings1051977-2 pandawani, Biodiversitas_Ni Putu Pandawani_2022_R1_REV_Complate (1).doc (2)	May 2, 2022	Research Results
Settings1051978-1 pandawani, Certificate # FP-2229.pdf	May 2, 2022	Other
Settings1051979-1 pandawani, Review Hermanu Triwidodo.doc	May 2, 2022	Other
Settings1051980-1 pandawani, Review Mimi Sutrawati.doc	May 2, 2022	Other

- [Download All Files](#)

Pre-Review Discussions

- [Add discussion](#)

Name	From	Last Reply	Replies	Closed
Settings Comments for the Editor	pandawani 2022-05-02 02:20 PM	-	0	<input type="checkbox"/>

Platform &
workflow by
OJS / PKP

Participants Edit

Smujo Editors (editors)

Ni Putu Pandawani (pandawani)

Messages

Note From

Dear Editor Team,

Our manuscript with the title "High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection". The high impact of pagoda leaf extract in suppressing Zucchini yellow mosaic virus (ZYMV) infection in zucchini plants was first reported in this study. The effect of application time and concentration of appropriate plant extracts in suppressing ZYMV infection is also a novelty in this study. Application of pagoda leaf extract one day before ZYMV inoculation with a concentration of 50% was able to reduce the incidence and severity of disease, accelerate flowering, and reduce yield loss of zucchini fruit. The results of this study enrich the information on the potential of pagoda flower leaves as agents for controlling plant viruses.

pandawani

2022-05-02 02:20 PM

My Assigned

Search

[New Submission](#)

Filters

Overdue

Incomplete

Stages

Submission

Review

Copyediting

Production

6

PANDAWANI et al. – *Effect of Clerodendrum paniculatum leaf extract on zucchini yellow mosaic virus infection*

Sections

Articles

Activity

Days since last activity

Days since last activity

[ID11097](#)

[Pandawani](#)

[High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection](#)

[Revisions have been requested.](#)

ReviewCurrently in the Review stage.

1/1 3

Hide expanded details about Pandawani

1/1 Assigned reviews completed

0 Revisions submitted

3 Open discussions

Last activity recorded on Wednesday, May 25, 2022.

Notifications^{undefined}

[biodiv] Editor Decision

2022-05-19 06:13 AM

Ni Putu Pandawani:

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection".

Our decision is: Revisions Required

Reviewer A:

Comments:

1. Some lines are confusing so please correct them.
2. To maintain the uniformity, i used pagoda extract instead of *C. paniculatum*.

3. Please check all the references again some are repeated twice.

4. correct the figure 2.

5. Please read paper very carefully.

Recommendation: See Comments

[undefined](#)

[biodiv] Editor Decision

2022-05-19 01:48 PM

Ni Putu Pandawani:

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection".

Our decision is: Revisions Required

Reviewer A:

Comments:

1. Some lines are confusing so please correct them.

2. To maintain the uniformity, i used pagoda extract instead of *C. paniculatum*.

3. Please check all the references again some are repeated twice.

4. correct the figure 2.

5. Please read paper very carefully.

Recommendation: See Comme

Notifications

[undefined](#)

[biodiv] Editor Decision

2022-05-24 03:04 PM

Ni Putu Pandawani:

We have reached a decision regarding your submission to Biodiversitas Journal of Biological Diversity, "High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection".

Our decision is: Revisions Required

Reviewer A:

Comments:

1. I have some corrections in your paper, so please check them.

2 read papervery carefully.

Recommendation: See Comments

11097 / **Pandawani** / High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection

Upload FileLibrary

WorkflowPublication

- [Submission](#)

- [Review](#)

- [Copyediting](#)

- [Production](#)

- [Round 1](#)

- [Round 2](#)

- [Round 3](#)

Round 1 Status New reviews have been submitted and are being considered by the editor.

Notifications

- [\[biodiv\] Editor Decision](#) 2022-05-19 06:13 AM
- [\[biodiv\] Editor Decision](#) 2022-05-19 01:48 PM
- [\[biodiv\] Editor Decision](#) 2022-05-24 03:04 PM

Reviewer's Attachments

- [Search](#)

Name	Date	Component
------	------	-----------

*No Files***Revisions**• [Search](#)

• [Upload File](#)

Name	Date	Component
------	------	-----------

*No Files***Review Discussions**• [Add discussion](#)

Name	From	Last Reply	Replies	Closed
Settings revision completed	pandawani 2022-05-20 10:17 AM	-	0	<input type="checkbox"/>
Uncorrected Proof	dewinurpratiwi 2022-05-25 04:38 AM	pandawani 2022-05-25 11:32 AM	2	<input type="checkbox"/>
BILLING	dewinurpratiwi 2022-05-25 04:50 AM	-	0	<input type="checkbox"/>

11097 / **Pandawani** / High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection
Upload FileLibrary
WorkflowPublication

- [Submission](#)

- [Review](#)

- [Copyediting](#)

- [Production](#)

- [Round 1](#)

- [Round 2](#)

- [Round 3](#)

Round 2 Status All reviewers have responded and a decision is needed.

Notifications

- [\[biodiv\] Editor Decision](#) 2022-05-19 06:13 AM
- [\[biodiv\] Editor Decision](#) 2022-05-19 01:48 PM
- [\[biodiv\] Editor Decision](#) 2022-05-24 03:04 PM

Reviewer's Attachments

- [Search](#)

Name	Date	Component
<input type="text" value="1052827-1"/> , 11097-Article Text-1052437-1-4-20220511.doc	May 19, 2022	

Revisions

• [Search](#)

• [Upload File](#)

Name	Date	Component
Settings¹⁰⁵²⁹²²⁻¹ Research Results, A-11097-Article Text-1052437-1-4-20220511_Rev1.doc	May 20, 2022	Research Results

Review Discussions

• [Add discussion](#)

Name	From	Last Reply	Replies	Closed
Settings revision completed	pandawani 2022-05-20 10:17 AM	-	0	<input type="checkbox"/>
Uncorrected Proof	dewinurpratiwi 2022-05-25 04:38 AM	pandawani 2022-05-25 11:32 AM	2	<input type="checkbox"/>
BILLING	dewinurpratiwi 2022-05-25 04:50 AM	-	0	<input type="checkbox"/>

Uncorrected Proof

[Close Panel](#)

Participants

- Smujo Editors (editors)
- Anisa Septiasari (aseptiasari)
- DEWI NUR PRATIWI (dewinurpratiwi)
- Ni Putu Pandawani (pandawani)

Messages

Note	From
<p>Dear Author(s),</p> <p>Pls, find attached file for an uncorrected proof (Copyedited file).</p> <p>The revised manuscript is awaited. Do not worry about layout changes due to revision; our staff will fix it again.</p> <p>Note: Kindly use track change when you make improvements.</p>	<p>dewinurpratiwi</p> <p>2022-05-25</p> <p>04:38 AM</p>

Note	From
dewinurpratiwi, Flowering - Pandawani.doc	
<p>Settings Dear Biodiversitas team, Thank you for your information. I have corrected the article according to suggestions from reviewers. can this article be included in the full article? not put into short communication. Best regards, Ni putu pandawani</p> <p>pandawani, A-11097-Research Results-1052925-1-4-20220520_R2.doc</p>	<p>pandawani 2022-05-25 11:21 AM</p>
<p>Settings Dear Biodiversitas team,</p> <p>I've checked and corrected the article in proofreading layout. Here I attach the article that I have corrected. Thank you very much</p> <p>Best regards,</p> <p>Ni Putu Pandawani</p> <p>pandawani, 11097-Article Text-1053214-1-18-20220525_Rev.doc</p>	<p>pandawani 2022-05-25 11:32 AM</p>
Add Message	

ARTIKEL.. SUBMITE REVISI .. 25 MEI 2022

High impact of pagoda leaf extract to suppress zucchini yellow mosaic virus infection in zucchini plants

¹Study Program of Agrotechnology, Faculty of Agriculture and Business, University of Mahasaraswati Denpasar, Bali 80233, Indonesia

²Study Program of Agrotechnology, Faculty of Agriculture, Warmadewa University, Bali 80239, Indonesia

[✉]email: pandawaniputu@unmas.ac.id

Abstract. Zucchini yellow mosaic virus (ZYMV) is an important virus in zucchini plants. ZYMV infection is prevalent on zucchini plants in Bali and is extremely difficult to eradicate. This study aimed to determine the efficacy of pagoda leaf extract in reducing yield loss in zucchini plants. A factorial randomized block design with two variables was employed in this study. The application time of the pagoda extract was the first variable, and concentration was the second. Plant extract was applied during seed immersion, one day before and after inoculation, and on the same day of virus inoculation. The concentrations of pagoda extract employed were 10%, 30%, and 50%. The crude extract was sprayed onto the zucchini leaves. Plants treated with extract one day after virus inoculation at 10%, 30%, and 50% concentrations exhibited yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting. While application of extract one day before virus inoculation at 30% and 50% concentrations resulted in yellowing and yellowing with cupping signs. Plants infected with ZYMV first showed flowering at 37 DAP, whereas plants treated with pagoda leaf extract one day before virus inoculation started flowering at 32 DAP. Application of pagoda leaf extract one day before ZYMV inoculation on zucchini plants effectively suppressed 40% yield loss. The findings of present investigation revealed information about the potential of pagoda leaf extract as a plant virus control agent through the mechanism of induction in zucchini plants. **Keywords:** Flowering, pagoda leaf extract, yield loss, zucchini, ZYMV

Abbreviations (if any): Zucchini yellow mosaic virus (ZYMV)

Running title: High impact of pagoda flower leaves to suppress zucchini yellow mosaic virus infection

INTRODUCTION

ZYMV is one of the most critical viruses causing mosaic disease in Cucurbitaceae plants worldwide, which could cause yield reduction (Simmons et al. 2011; Simmons et al. 2013). The incidence of viral disease due to ZYMV infection in cucumber plants could reach 100%, with varying attack intensities

(Tymchyshyn et al. 2017). The loss due to viral infection depends on the time of infection and can result in yield losses of up to 100% (Coutts et al. 2011). Plants become very susceptible to viral infection at young age, affecting the high incidence of disease. When young plants become infected, the incubation period of virus is shorter, and the virus distribution and translocation process is accelerated (Coutts et al. 2013). Host range is one way of determining the biological nature of a virus (Spadotti et al. 2015). The most frequently employed host range test plants for ZYMV are *Gomphrena globosa* L., *Chenopodium amaranticolor* Coste et Reyn, *C. quinoa* L., *Citrullus lanatus*, *Cucumis melo* L., *Cucumis sativus* L., *Luffa acutangula*, *Momordica charantia*, *Sechium edule*, *Phaseolus vulgaris* L., *Vigna sinensis*, *Capsicum annuum*, *Datura stramonium*, *Lycopersicon esculentum*, *Nicotiana tabacum* L. cv. White barley, *N. benthamiana*, and *Physalis floridana* (Jaroszewska et al. 2013; Massumi et al. 2013; Spadotti et al. 2015).

Zucchini yellow mosaic virus (ZYMV) has a one-molecule positive senseRNA genome of approximately 10 kb and encodes a poly-protein that is proteolytically processed into mature protein; P1 (protease), HC (helper component/protease), P3, 6K1, CI (cylinder inclusion), 6K2, NIa (nuclear inclusion a), VPg (viral protein-associated genome), NIb (nuclear inclusion b) and CP (Gal-On 2007; Moradi et al. 2019). Until now, four out of ten potyviral proteins have been identified in ZYMV-infected plants, namely P1, HC-Pro, cylindrical inclusions, cylindrical inclusions (CI) pinwheel type 1, and CP (Maghamnia et al. 2018; Valli et al. 2017). Other proteins (Third protein (P3), 6K1, 6K2, viral genome-linked protein (VPg), Nuclear Inclusion A (NIa), and NIb replicas) have not been identified in ZYMV-infected plants and are based on potyvirus sequence homology.

ZYMV was first reported in Italy in 1973 (Lisa et al. 1981) and then France (Lecoq et al. 1981). ZYMV distribution has been found in Africa, America, Asia, Europe, the Middle East, and Oceania (Spadotti et al. 2015; Bubici et al. 2020). In Egypt, ZYMV has been associated with significant yield losses in many cucurbit crops, including cucumbers, melons, pumpkins, marrow vegetables, watermelons, and especially zucchini (Aleem et al. 2021). ZYMV infection result in symptoms, such as mosaicism, yellowing, and eventually "shoeing" on the leaves. The fruit is stunted, twisted, and distorted, resulting in reduced yield and inability to market the product, especially chayote zucchini (Massumi et al. 2011). ZYMV is commonly transferred in two ways: horizontally via aphid vectors and vertically by transmission from the first generation of ZYMV-infected seeds to the following generation of ZYMV-infected seeds (Simmons et al. 2011; Wang and Li 2017). Several species of aphids are ZYMV vectors that transmit viruses non-persistently, such as *A. gossypii*, *Myzus persicae* (Maina et al. 2017; Romay et al. 2014), *Asyrthosiphon pisum*, *A. kondoi*, *Aphis craccivora*, *A. citricola*, *A. middletonii*, *A. spiraecola*, *Macrosiphum euphorbiae*, *Toxoptera aurantii*, and *Uroleucon ambrosiae* (Spadotti et al. 2015). ZYMV can also be transmitted mechanically easily through cutting tools that have been contaminated with the virus (Nasr-Eldin et al. 2016; Prendeville et al. 2012).

Viral diseases can be controlled by eradicating insect vectors, quarantining, and planting healthy and virus-free seeds. Additionally, the usage of antiviral compounds derived from plant extracts has been proven to be effective against a variety of viruses due to their ribosome-inactivating proteins (RIPs) (Gal-On 2007). One of the agents capable of inducing systemic resistance in plants is a plant extract (Elbeshehy 2017). The effective control of viral diseases is the use of resistant varieties, but there are not many commercial virus-resistant cultivars available.

Systemic resistance of a plant can be activated by inducing resistance genes present in plants by utilizing resistance-inducing agents (Elbeshehy 2017). One of the agents that induce systemic resistance of plants is plant extracts (Gal-On 2007). Spinach leaf extract (*Amaranthus spinosus*), four o'clock flower leaf (*Mirabilis jalapa*), and pagoda leaf (*Clerodendrum paniculatum*) were reported to induce systemic resistance to anthracnose and CMV pathogens in chili (Hersanti 2003). Other plant extract such as *Chenopodium amaranticolor* is reported to have antiviral activity for the tobacco mosaic virus (TMV) and antitumor for Ehrlich tumor (EA) (De Oliveira et al. 1993). Extracts of pagoda leaf, thorn spinach, four o'clock flower, *C. amaranticolor*, and sambiloto are able to suppress BCMV infection in yardlong bean plants (Kurnianingsih and Damayanti 2012).

The utilization of efficacious plant extracts have been used to benefit human health, but are still of little use in plant protection. Therefore, it is necessary to explore potential plant species to control pathogens, including viruses. The objective of this study was to test the effectiveness of plant extract to suppress ZYM

material and METHODS

Preparation of zucchini plant (*Cucurbita pepo* L.)

The seeds of the Carisa cultivar zucchini were sown on plastic composite trays containing sterile growth media of a mixture of soil and compost. After 15 days, the grown seedlings were transplanted into plastic pots measuring 35 cm × 30 cm containing 5 kg of sterile planting medium, i.e. soil and manure (2:1 ratio). Each pot was planted with three seeds, and after one week of growth, one healthy plant was selected for the research object. Plants were irrigated daily with appropriate volume.

Propagation of ZYMV inoculum

The ZYMV inoculum was obtained from the Laboratory of the Faculty of Agriculture, Mahasaraswati University in Denpasar, then confirmed its nucleic acid sequence via DNA sequencing and designated it as ZYMV-zucchini isolate Bali (Pandawani and Widnyana 2021) (Figure 1). The inoculum was propagated mechanically by inoculating ZYMV inoculum on zucchini plants two weeks after planting. Mechanical transmission of ZYMV was carried out by grinding sick leaves (source of inoculum) in phosphate buffer pH 7 with a ratio of 1:10 (w/v). The sap was then applied to the leaves of healthy plants which were treated with 600 mesh carborundum. After inoculation, leaves were rinsed with running water. The plants were then reared in a greenhouse, and symptomatic young leaves were used as a source of inoculum in this study.

Preparation and application of pagoda (*Clerodendrum Paniculatum*) extract

The leaves of pagoda plant (*C. paniculatum*) was obtained from the area around Baturiti Tabanan. Pagoda extract was prepared through a drying process to obtain a simplicia form. The simplicia was then macerated to produce a concentrated leaf extract with a concentration of 100% (Verma et al. 1998). Application of pagoda extract to zucchini plants was carried out according to the application time determined with a volume of 100 mL per plant. The pagoda extract was applied on zucchini leaves after three weeks of transplanting.

Inoculation of ZYMV in zucchini plants (*Cucurbita pepo* L.)

Three-week-old zucchini plants were mechanically inoculated with ZYMV. Virus inoculum was prepared by grinding symptomatic young leaves of zucchini plant (source of ZYMV inoculum) in 1:10 (w/v) phosphate buffer. The virus was inoculated by applying the inoculum on the first leaf of zucchini plant, which was sprinkled with 600 mesh carborundum.

Experiment design and data analysis

This study was conducted in a completely randomized design with a factorial pattern. The first factor was the time of application of the inducer extract, which consisted of five treatments, namely: 1) application of pagoda extract during seed soaking (T1), 2) application time of pagoda extract one day before inoculation of ZYMV sap (T2), 3) application of pagoda extract on the same day as ZYMV applied (T3), 4) application of pagoda extract one day after ZYMV (T4), and no application of pagoda extract (T0) (control). The second factor was the concentration of inducer extract, which consisted three treatments, namely: 1) 10% of pagoda extract (C1), 2) 30% of pagoda extract (C2), and 3) 50% of pagoda extract (C3). The treatments were combined to obtain (5 × 3) 15 combination treatments, and each combination treatment was repeated ten times to obtain 45 experimental units. Observation parameters included disease incidence, type of disease symptoms, disease severity, virus concentration, and loss of fruit yield. Data were analyzed using ANOVA according to a completely randomized design with a factorial pattern (5 × 3 × 10) with Duncan's follow-up test (DMRT) at 5% level.

Enzyme-linked immunosorbent assays (ELISA)

The detection of ZYMV virus was performed on the leaves of zucchini plant four weeks after inoculation. Serological virus detection was carried out by enzyme-linked immunosorbent assay (DAS-ELISA) using Potyvirus antiserum (DSMZ). Virus accumulation was quantitatively read using ELISA Reader model 550 (Bio-Rad, USA) at a wavelength of 405 nm.

RESULTS AND DISCUSSION

Incubation period

The incubation period of ZYMV in plants treated with pagoda leaf extract was the most prolonged average incubation period (18.27 DAP) compared to the treatment without pagoda leaf extract (C+) (3.60 DAP). Both factors (inoculation time and concentration of pagoda leaf extract) exhibited a significantly longer incubation period when compared with the control treatment. The application of 50% pagoda leaf extract one day before virus inoculation significantly slowed the incubation period of virus (21.00 DAP) compared with other treatments (Table 1). Kurnianingsih and Damayanti (2012) found that pagoda leaf extract could slow the incubation period of BCMV in yardlong beans.

Symptoms of infection

ZYMV inoculation produced various symptoms, such as yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting (Figure 2 and Table 1). Plants treated with leaf extract one day after virus inoculation at concentrations of 10%, 30%, and 50% showed symptoms of yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting. The application of extract one day before inoculation of viral sap at concentrations of 30% and 50% only produced yellowing and yellowing with cupping symptoms.

Disease incidence

Plants treated with pagoda leaf extract one day before sap virus inoculation showed the lowest (20%) disease incidence until the end of observation. The other treatments resulted in 40% - 60% disease prevalence. In control plants where no pagoda leaf extract was applied, disease incidence reached 100% (Table 4).

Disease severity

Pagoda extract treatment significantly reduced the severity of disease. Among the treatments, the highest disease severity was recorded in extract application treatment one day after inoculation of viral smear. In comparison, the lowest was observed in the application of extract one day before the inoculation of virus (Table 2). The best time for application of pagoda leaf extract was one day before ZYMV inoculation with a concentration of 50% because it could reduce disease severity up to 60.79% (Table 4). It was observed that pagoda leaf extract suppressed the disease incidence and severity of ZYMV infection. This is in accordance with Damayanti and Pebriyeni (2015), Kurnianingsih and Damayanti (2012), who reported that pagoda leaf extract can suppress BCMV infection reaching 100% in yardlong beans.

Virus detection with ELISA

ELISA absorbance value (NAE) showed that no virus was detected in the samples given pagoda flower leaf extract at a concentration of 30% and 50% one day before virus inoculation. NAE was not significantly different from the negative control (Table 4). Based on ELISA test, zucchini plants showed presence (positive) of ZYMV in all treatments, except one day before virus sap inoculation with a concentration of 30 and 50%, zucchini plants showed absence (negative) of ZYMV.

Flowering and yield loss

Infection with ZYMV inhibits flowering in zucchini plants. ZYMV-infected plants showed reduced flowering and yield loss in zucchini (Kheder et al. 2017; Kone et al. 2017; Maghamnia et al. 2018; Maina et al. 2017). Plants infected with ZYMV first flowered at 37 DAP, while plants treated with pagoda leaf extract one day before virus inoculation was flowered at 32 DAP (Table 3). Treatment of pagoda leaf extract one day before virus inoculation on zucchini plants resulted in a yield loss of 19.99% - 23.50% compared with other treatments. This indicated that applying pagoda leaf extracts one day before ZYMV inoculation on zucchini plants effectively reduced yield loss by 40%. Application of pagoda leaf extract one day before virus inoculation increased the resistance of zucchini plants. The low severity and virus accumulation in this treatment indicated the potential of the pagoda leaf as an inducer of systemic resistance of zucchini plants. The mechanism of induction of systemic resistance of zucchini by pagoda leaf extract has been reported by Verma et al. (1998). The active compound in the form of 34 kDa protein in pagoda flower leaf extract causes tobacco leaves to become immune to viruses. In addition, pagoda flower leaf extract can suppress BCMV because it contains virus inhibitors and has antiviral activity (De Oliveira et al. 1993). The active compounds of the pagoda flower are known as antiviral proteins and are known as ribosome-inactivating proteins (RIPs). RIPs are also present in the root and leaf extract of *M.*

jalapa and are referred to as Mirabilis antiviral protein (MAP) (Verma et al. 1998). MAP can reach the ribosome's active site earlier than the virus to prevent viral infection at an early stage before the virus undergoes encapsidation (Vivanco et al. 1999).

In the present study pagoda leaf extract showed 80% effectiveness in inhibiting ZYMV infection, whereas against BCMV and CMV, inhibition was observed to be 100% and 82.6%, respectively (Hersanti 2003; Kurnianingsih and Damayanti 2012). It indicates that the same plant extract showed different effectiveness against different viruses.

Application of pagoda leaf extract one day before ZYMV inoculation with a concentration of 50% was able to reduce disease incidence and severity of, accelerate flowering, and reduce yield loss of zucchini fruit compared to the control treatment. The results of this study enrich the information about the potential of pagoda leaves as agent to control plant viruses.

ACKNOWLEDGEMENTS

Acknowledgments are conveyed to the farmer groups in Banjar Sandan Village, Baturiti, Tabanan, Bali for their assistance and support.

REFERENCES

- Aleem EEA, Rabie M, Fattouh FA. 2021. Molecular characterisation of zucchini yellow mosaic virus infecting *Cucurbita pepo* in Egypt. Plant Prot Sci. DOI: 10.17221/191/2020-PPS
- Bubici G, Navarro B, Carluccio AV, Ciuffo M, Serio FD, Fabrizio Cillo F. 2020. Genomic sequence variability of an Italian zucchini yellow mosaic virus isolate. Eur J Plant Pathol 156: 325-332.
- Coutts BA, Kehoe MA, Jones RAC. 2011. Minimising losses caused by zucchini yellow mosaic virus in vegetable cucurbit crops in tropical, subtropical and Mediterranean environments through cultural methods and host resistance. Virus Res. DOI: 10.1016/j.virusres.2011.04.015
- Coutts BA, Kehoe MAA, Jones RA. 2013. Zucchini yellow mosaic virus: Contact transmission, stability on surfaces, and inactivation with disinfectants. Plant Dis. DOI: 10.1094/PDIS-08-12-0769-RE
- Damayanti TA, Pebriyeni L. 2015. Barrier crop and pagoda leaf extract to control Bean common mosaic virus on yard long bean in the field. J Hort. 25: 238-245.

- De Oliveira MM, Sampaio MR, Noronha AB, Vicente M, Vianna S. 1993. Detection of antiviral and antitumoral fractions of *Chenopodium amaranticolor* leaf extract. *Microbios* 76: 213-221.
- Elbeshehy EF. 2017. Inhibitor activity of different medicinal plants extracts from *Thuja orientalis*, *Nigella sativa* L., *Azadirachta indica* and *Bougainvillea spectabilis* against Zucchini yellow mosaic virus (ZYMV) infecting *Citrullus lanatus*. *Biotechnol Biotechnol Equip*. DOI: 10.1080/13102818.2017.1279572
- Gal-On A. 2007. Zucchini yellow mosaic virus: Insect transmission and pathogenicity-The tails of two proteins. *Mol Plant Pathol*. DOI: 10.1111/j.1364-3703.2007.00381.x
- Hersanti. 2003. Testing the potential of extracts of 37 plant species as agents of inducing systemic resistance of red chili plants to Cucumber mosaic virus. *J Phytopathol Indones* 7: 54-58.
- Jaroszewska BH, Rymelska N, Borodynko N, Pospieszny H. 2013. Biological and molecular characterization of the Polish zucchini yellow mosaic virus isolates. *Acta Sci Pol Hortorum Cultus* 12: 75-85.
- Kheder AA, Taqwa M, Sulaiman TM, Ghanem GAM, Tohamy MR. 2017. Biological, serological and molecular characterization of Egyptian zucchini yellow mosaic virus isolate infecting squash plants in Fayoum governorate. *Egypt J Phytopathol*. DOI: 10.21608/ejp.2017.88570
- Kone N, Asare-Bediako E, Silue S, Kone D, Koita O, Wulf M, Stephan W. 2017. Influence of planting date on incidence and severity of viral disease on cucurbits under field condition. *Ann Agric Sci*. DOI: 10.1016/j.aogas.2017.05.005
- Kurnianingsih L, Damayanti TA. 2012. Five plant extracts for suppressing Bean common mosaic virus infection on yard long bean. *J Phytopathol Indones*. DOI: 10.14692/jfi.8.6.155
- Lecoq H, Pitrat M, Cle'ment M. 1981. Identification et caracterisation d'un potyvirus provoquant la maladie du rabougrissement jaune du melon. *Agronomie* 1: 827-834.
- Lisa V, Boccardo G, D'Agostino G, Dellavalle G, D'Aquilio M. 1981. Characterization of a potyvirus that causes zucchini yellow mosaic. *Phytopathol* 71: 667-672.
- Maghamnia HR, Hajizadeh M, Azizi A. 2018. Complete genome sequence of zucchini yellow mosaic virus strain Kurdistan, Iran. *3 Biotech*. DOI: 10.1007/s13205-018-1177-3

- Maina S, Coutts BA, Edwards OR, Almeida L, Kehoe MA, Ximenes A, Jones RAC. 2017. Zucchini yellow mosaic virus populations from East Timorese and Northern Australian cucurbit crops: Molecular properties, genetic connectivity, and biosecurity implications. *Plant Dis*. DOI: 10.1094/PDIS-11-16-1672-RE
- Massumi H, Shaabani M, Heydarnejad J, Hosseini Pour AH, Rahimian H. 2011. Host range and phylogenetic analysis of Iranian isolates of zucchini yellow mosaic virus. *J Plant Pathol* 93: 187-193.
- Moradi Z, Mehrvar M, Nazifi E. 2019. Population genetic analysis of zucchini yellow mosaic virus based on the CI gene sequence. *J Cell Mol Res*. DOI: 10.22067/jcmr.v10i2.76133
- Nasr-Eldin MA, Hayam S, Abdelkader HS, Abo-Senna AS, Othman BA. 2016. Characterization and phylogenetic analysis of zucchini yellow mosaic virus infecting *Cucurbita pepo* in Egypt. *Am J Sci*. DOI: 10.7537/marsjas12031613
- Pandawani NP, Widnyana IK. 2021. Identification of virus causes of mosaic diseases in zucchini plants in the Bali Island of Indonesia. *J Tekirdag Agric Fac*. DOI: 10.33462/jotaf.707645
- Prendeville HR, Ye X, Morris J, Pilson D. 2012. Virus infections in wild plants are both frequent and often unapparent. *Am J Bot*. DOI: 10.3732/ajb.1100509
- Romay G, Lecoq H, Geraud-Pouey F, Chirinos DT, Desbiez C. 2014. Current status of cucurbit viruses in Venezuela and characterization of Venezuelan isolates of zucchini yellow mosaic virus. *Plant Pathol*. DOI: 10.1111/ppa.12072
- Simmons HE, Dunham JP, Zinn KE, Munkvold GP, Holmes EC, Stephenson AG. 2013. Zucchini yellow mosaic virus (ZYMV, Potyvirus): Vertical transmission, seed infection and cryptic infections. *Virus Res*. DOI: 10.1016/j.virusres.2013.06.016
- Simmons HE, Holmes EC, Gildow FE, Bothe-Goralczyk MA, Stephenson AG. 2011. Experimental verification of seed transmission of zucchini yellow mosaic virus. *Plant Dis*. DOI: 10.1094/PDIS-11-10-0843
- Spadotti D, Wassano D, Rezende J, Camargo LEA, Inoue-Nagata AK. 2015. Biological and molecular characterization of Brazilian isolates of zucchini yellow mosaic virus. *Sci Agric*. DOI: 10.1590/0103-9016-2014-0197
- Tymchyshyn O, Shevchenko T, Shevchenko O, Budzanivska I. 2017. Phylogenetic analysis of seed-transmitted isolate of zucchini yellow mosaic virus. *Bulletin of Taras Schevchenko National University of Kyiv* 2: 47-50.
- Valli AA, Gallo A, Rodamilans B, López-Moya JJ, García JA. 2017. The HCPro from the Potyviridae family: an enviable multitasking helper component that every virus would like to have. *Mol Plant Pathol*. DOI: 10.1111/mpp.12553

Verma HN, Baranwal VK, Srisavasta S. 1998. Antiviral substances of plant origin. In: Hadidi A, Khetarpal RK, Koganezawa H, editor. *Plant Virus Disease Control*. St. Paul Minnesota (US): APS Pr. Page 154-162.

Vivanco JM, Querci M, Salazar LF. 1999. Antiviral and antiviroid activity of MAP containing extracts from *Mirabilis jalapa* roots. *Plant Dis*. DOI: 10.1094/PDIS.1999.83.12.1116

Wang D, Li G. 2017. Biological and molecular characterization of zucchini yellow mosaic virus isolates infecting melon in Xinjiang, China. *Can J Plant Pathol*. DOI: 10.1080/07060661.2017.1285356

TABLES LIST

Table 1. Incubation period in ZYMV-infected zucchini plants in response to differences in application time and concentration of pagoda leaf extract (*C. paniculatum*)

Pagoda leaf extract concentration (%)	Incubation period of ZYMV infection (DAP)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	3.40 a	14.0 e	16.00 fg	6.20 b	6.00 b	9.12 a
30	3.60 a	15.2 f	17.80 h	8.60 d	7.20 c	10.48 b
50	3.80 a	16.8 g	21.00 i	9.40 d	8.40 d	11.88 c

Average of extract concentration	3.60 a	15.33 d	18.27 e	8.07 c	7.20 b
----------------------------------	--------	---------	---------	--------	--------

Note: day after planting (DAP); Different letters in the same row show a significant effect compared to the positive control (C+)

Table 2. Disease severity in ZYMV-infected zucchini plants in response to differences in application time and concentration of pagoda leaf extract (*C. paniculatum*)

Pagoda leaf extract concentration (%)	Disease severity (%)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	66.37 j	39.96 e	31.80 bc	51.42 g	61.93 hij	50.30 c
30	64.30 ij	37.01 cd	27.33 ab	49.71 fg	58.78 hi	47.43 b
50	62.59 j	32.87 bc	23.59 a	45.14 ef	56.88 h	44.21 a
Average of extract	64.42 e	36.61 b	27.57 a	48.76 c	59.19 d	

concentration

Note: Different letters in the same row show a significant effect compared to the positive control (C+)

Table 3. Flowering in zucchini plants in response to differences in application time and concentration of pagoda leaf extract

Pagoda leaf extract concentration (%)	Appearance of flower (DAP)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	37.00 f	33.60 bcd	32.80 b	35.80 e	36.80 f	35.20 b
30	37.20 f	32.60 b	33.80 bcd	33.20 bc	37.40 f	34.84 b
50	36.80 f	33.40 bc	30.80 a	34.60 cd	35.00 de	34.12 a
Average of extract concentration	37.00 c	33.20 a	32.47 a	34.53 b	36.40 c	

Note: day after planting (DAP); Different letters in the same row show a significant effect compared to the positive control (C+)

Table 4. Effect of application time and concentration of pagoda leaf extract (*C. paniculatum*) on disease incidence, type of symptoms, suppression of ZYMV disease severity, and yield loss

Application time of pagoda leaf extract	Extract concentration (%)	Disease incidence (%)	Type of disease symptom*	Disease severity suppression (%)	ELISA results		Yield loss (%)
					Average of NAE**	Reaction	
C+	10	100 (10/10)	M, Mc, St	-	0.418	+	58.42
	30	100 (10/10)	M, Mc, St	-	0.43	+	55.90
	50	60 (6/10)	Y, Yc, M	-	0.395	+	55.29
Extract application during seed soaking	10	60 (6/10)	M, Mc, St	39.77	0.557	+	35.56
	30	40 (4/10)	Y, Yc, Mc, St	42.58	0.486	+	33.69
	50	40 (4/10)	Y, Yc, Mc, St	47.70	0.458	+	37.41
Extract application one day before virus sap inoculation	10	40 (4/10)	Y, Yc	52.04	0.426	+	23.50
	30	20 (2/10)	Y, Yc	58.99	0.184	-	22.18
	50	20 (2/10)	Y, Yc	60.79	0.112	-	19.99

Extract application concurrently with viral sap inoculation	10	60 (6/10)	M, Mc, St	22.57	0.608	+	55.31
	30	40 (4/10)	M	23.25	0.515	+	40.74
	50	40 (4/10)	Y, Yc	26.96	0.489	+	43.93
Extract application one day after viral sap inoculation	10	60 (6/10)	Y, Yc, M, Mc, St	6.68	0.638	+	55.44
	30	60 (6/10)	M	8.96	0.588	+	55.19
	50	40 (4/10)	M	8.90	0.550	+	54.63

Note: C+ (without extract application)

Type of disease symptom*: Y = yellowing, Yc = yellowing with cupping, M = mosaic, Mc = mosaic with cupping, St = stunting

Negative control: healthy plants (without virus inoculation and extract application)

Positive control: sick plants (virus inoculation and without extract application)

**NAE: Elisa Absorbance value

FIGURES LIST



Figure 1. Inoculum source of ZYMV isolates Bali on zucchini plants




Figure 2. Disease symptoms of ZYMV isolates Bali on zucchini plants: A. mosaic, B. yellowing with cupping, C. mosaic with cupping, and D. yellowing

Dear Author(s),

Pls, find attached file for an uncorrected proof (Copyedited file).

The revised manuscript is awaited. Do not worry about layout changes due to

dewinurpratiwi

2022-05-25 04:38

AM

revision; our staff will fix it again.

Note: Kindly use track change when you make improvements.

[dewinurpratiwi, Flowering - Pandawani.doc](#)

Short Communication:

High impact of *Clerodendrum paniculatum* leaf extract to suppress zucchini yellow mosaic virus infection in zucchini plants

NI PUTU PANDAWANI^{1,✉}, LISTIHANI¹, I KETUT WIDNYANA¹, PUTU EKA PASMIDI ARIATI¹, DEWA GEDE WIRYANGGA SELANGGA²

¹Program of Agrotechnology, Faculty of Agriculture and Business, Universitas Mahasaraswati Denpasar. Jl. Kamboja No.11A, Dangin Puri Kangin, Kec. Denpasar Utara, Kota Denpasar, Bali 80233. Tel./fax. +62- 361-227019, ✉email: pandawaniputu@unmas.ac.id

²Program of Agrotechnology, Faculty of Agriculture, Universitas Warmadewa. Jl. Terompong No.24, Sumerta Kelod, Kec. Denpasar Tim., Kota Denpasar, Bali 80239

Manuscript received: 2 May 2022. Revision accepted: xxx May 2022.

Abstract. Pandawani NP, Listihani, Wduama IK, Ariati PEP, Selangga DGW. 2022. Short Communication: High impact of *Clerodendrum paniculatum* leaf extract to suppress zucchini yellow mosaic virus infection in zucchini plants. *Biodiversitas* 23: xxxx. Zucchini yellow mosaic virus (ZYMV) is an important virus in zucchini plants (*Cucurbita pepo* L.). ZYMV infection is prevalent on *C. pepo* in Bali and is extremely difficult to eradicate. This study aimed to determine the efficacy of pagoda leaf extract (*Clerodendrum paniculatum* L.) in reducing yield loss in *C. pepo*. A factorial randomized block design with two variables was employed in this study. The application time of the *C. paniculatum* extract was the first variable, and concentration was the second. Plant extract was applied during seed immersion, one day before and after inoculation, and on the same day of virus inoculation. The concentrations of *C. paniculatum* extract employed were 10%, 30%, and 50%. The crude extract was sprayed onto the *C. pepo* leaves. Plants treated with extract one day after virus inoculation at 10%, 30%, and 50% concentrations exhibited yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting. While application of extract one day before virus inoculation at 30% and 50% concentrations resulted in yellowing and yellowing with cupping signs. Plants infected with ZYMV first showed flowering at 37 DAP, whereas plants treated with *C. paniculatum* leaf extract one day before virus inoculation started flowering at 32

DAP. Application of *C. paniculatum* leaf extract one day before ZYMV inoculation on *C. pepo* effectively suppressed 40% yield loss. The findings of present investigation revealed information about the potential of *C. paniculatum* leaf extract as a plant virus control agent through the mechanism of induction in *C. pepo*.

Keywords: Flowering, pagoda leaf extract, yield loss, zucchini, ZYMV

Abbreviations: Zucchini yellow mosaic virus (ZYMV)

External



Ahmad Dwi Setyawan <smujo.id@gmail.com>

to me

Ni Putu Pandawani:

Thank you for submitting the manuscript, "High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection" to Biodiversitas Journal of Biological Diversity. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Submission URL: <https://smujo.id/biodiv/authorDashboard/submission/11071>

Username: pandawani

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Ahmad Dwi Setyawan

INTRODUCTION

ZYMV is one of the most critical viruses causing mosaic disease in Cucurbitaceae plants worldwide, which could cause yield reduction (Simmons et al. 2011; Simmons et al. 2013). The incidence of viral disease due to ZYMV infection in cucumber plants could reach 100%, with varying attack intensities (Tymchyshyn et al. 2017). The loss due to viral infection depends on the time of infection and can result in yield losses of up to 100% (Coutts et al. 2011). Plants become very susceptible to viral infection at young age, affecting the high incidence of disease. When young plants become infected, the incubation period of virus is shorter, and the virus distribution and translocation process is accelerated (Coutts et al. 2013). Host range is one way of determining the biological nature of a virus (Spadotti et al. 2015). The most frequently employed host range test plants for ZYMV are *Gomphrena globosa*

L., *Chenopodium amaranticolor* Coste et Reyn, *C. quinoa* L., *Citrullus lanatus*, *Cucumis melo* L., *Cucumis sativus* L., *Luffa acutangula*, *Momordica charantia*, *Sechium edule*, *Phaseolus vulgaris* L., *Vigna sinensis*, *Capsicum annum*, *Datura stramonium*, *Lycopersicon esculentum*, *Nicotiana tabacum* L. cv. White barley, *N. benthamiana*, and *Physalis floridana* (Jaroszewska et al. 2013; Massumi et al. 2013; Spadotti et al. 2015).

Zucchini yellow mosaic virus (ZYMV) has a one-molecule positive senseRNA genome of approximately 10 kb and encodes a poly-protein that is proteolytically processed into mature protein; P1 (protease), HC (helper component/protease), P3, 6K1, CI (cylinder inclusion), 6K2, NIa (nuclear inclusion a), VPg (viral protein-associated genome), NIB (nuclear inclusion b) and CP (Gal-On 2007; Moradi et al. 2019). Until now, four out of ten potyviral proteins have been identified in ZYMV-

infected plants, namely P1, HC-Pro, cylindrical inclusions, cylindrical inclusions (CI) pinwheel type 1, and CP (Valli et al. 2017; Maghamnia et al. 2018). Other proteins (Third protein (P3), 6K1, 6K2, viral genome-linked protein (VPg), Nuclear Inclusion A (NIa), and NIb replicas) have not been identified in ZYMV-infected plants and are based on potyvirus sequence homology.

ZYMV was first reported in Italy in 1973 (Lisa et al. 1981) and then France (Lecoq et al. 1981). ZYMV distribution has been found in Africa, America, Asia, Europe, the Middle East, and Oceania (Spadotti et al. 2015; Bubici et al. 2020). In Egypt, ZYMV has been associated with significant yield losses in many cucurbit crops, including cucumbers, melons, pumpkins, marrow vegetables, watermelons, and especially zucchini (*Cucurbita pepo* L.) (Aleem et al. 2021). ZYMV infection result in symptoms, such as mosaicism, yellowing, and eventually "shoeing" on the leaves. The fruit is stunted, twisted, and distorted, resulting in reduced yield and inability to market the product, especially chayote zucchini (Massumi et al. 2011). ZYMV is commonly transferred in two ways: horizontally via aphid vectors and vertically by transmission from the first generation of ZYMV-infected seeds to the following generation of ZYMV-infected seeds (Simmons et al. 2011; Wang and Li 2017). Several species of aphids are ZYMV vectors that transmit viruses non-persistently, such as *A. gossypii*, *Myzus persicae* (Maina et al. 2017; Romay et al. 2014), *Asyrthosiphon pisum*, *A. kondoi*, *Aphis craccivora*, *A. citricola*, *A. middletonii*, *A. spiraecola*, *Macrosiphum euphorbiae*, *Toxoptera aurantii*, and *Uroleucon ambrosiae* (Spadotti et al. 2015). ZYMV can also be transmitted mechanically easily through cutting tools that have been contaminated with the virus (Nasr-Eldin et al. 2016; Prendeville et al. 2012).

Viral diseases can be controlled by eradicating insect vectors, quarantining, and planting healthy and virus-free seeds. Additionally, the usage of antiviral compounds derived from plant extracts has been proven to be effective against a variety of viruses due

to their ribosome-inactivating proteins (RIPs) (Gal-On 2007). One of the agents capable of inducing systemic resistance in plants is a plant extract (Elbeshehy 2017). The effective control of viral diseases is the use of resistant varieties, but there are not many commercial virus-resistant cultivars available.

Systemic resistance of a plant can be activated by inducing resistance genes present in plants by utilizing resistance-inducing agents (Elbeshehy 2017). One of the agents that induce systemic resistance of plants is plant extracts (Gal-On 2007). Spinach leaf extract (*Amaranthus spinosus*), four o'clock flower leaf (*Mirabilis jalapa*), and pagoda leaf (*Clerodendrum paniculatum* L.) were reported to induce systemic resistance to anthracnose and CMV pathogens in chili (Hersanti 2003). Other plant extract such as *Chenopodium amaranticolor* is reported to have antiviral activity for the tobacco mosaic virus (TMV) and antitumor for Ehrlich tumor (EA) (De Oliveira et al. 1993). Extracts of *C. paniculatum* leaf, thorn spinach, four o'clock flower, *C. amaranticolor*, and sambiloto are able to suppress BCMV infection in yardlong bean plants (Kurnianingsih and Damayanti 2012).

The utilization of efficacious plant extracts have been used to benefit human health, but are still of little use in plant protection. Therefore, it is necessary to explore potential plant species to control pathogens, including viruses. The objective of this study was to test the effectiveness of plant extract to suppress ZYMV.

materials and METHODS

Preparation of zucchini plant (*Cucurbita pepo* L.)

The seeds of the Carisa cultivar *C. pepo* were sown on plastic composite trays containing sterile growth media of a mixture of soil and compost. After 15 days, the grown seedlings were transplanted into plastic pots measuring 35 cm × 30 cm containing 5 kg of sterile planting medium, i.e. soil and manure (2:1 ratio). Each pot was planted with three seeds, and

after one week of growth, one healthy plant was selected for the research object. Plants were irrigated daily with appropriate volume.

Propagation of ZYMV inoculum

The ZYMV inoculum was obtained from the Laboratory of the Faculty of Agriculture, Mahasaraswati University in Denpasar, then confirmed its nucleic acid sequence via DNA sequencing and designated it as ZYMV-zucchini isolate Bali (Pandawani and Widnyana 2021) (Figure 1). The inoculum was propagated mechanically by inoculating ZYMV inoculum on *C. pepo* two weeks after planting. Mechanical transmission of ZYMV was carried out by grinding sick leaves (source of inoculum) in phosphate buffer pH 7 with a ratio of 1:10 (w/v). The sap was then applied to the leaves of healthy plants which were treated with 600 mesh carborundum. After inoculation, leaves were rinsed with running water. The plants were then reared in a greenhouse, and symptomatic young leaves were used as a source of inoculum in this study.

Preparation and application of *Clerodendrum paniculatum* extract

The leaves of *C. paniculatum* was obtained from the area around Baturiti, Tabanan, Bali, Indonesia. *C. paniculatum* extract was prepared through a drying process to obtain a simplicia form. The simplicia was then macerated to produce a concentrated leaf extract with a concentration of 100% (Verma et al. 1998). Application of *C. paniculatum* extract to *C. pepo* was carried out according to the application time determined with a volume of 100 mL per plant. The *C. paniculatum* extract was applied on *C. pepo* leaves after three weeks of transplanting.

Inoculation of ZYMV in zucchini plants (*Cucurbita pepo* L.)

Three-week-old *C. pepo* were mechanically inoculated with ZYMV. Virus inoculum was prepared by grinding symptomatic young leaves of *C. pepo* plant (source of ZYMV inoculum) in 1:10 (w/v) phosphate buffer. The virus was inoculated by

applying the inoculum on the first leaf of *C. pepo* plant, which was sprinkled with 600 mesh carborundum.

Experiment design and data analysis

This study was conducted in a completely randomized design with a factorial pattern. The first factor was the time of application of the inducer extract, which consisted of five treatments, namely: (i) application of *C. paniculatum* extract during seed soaking (T1), (ii) application time of *C. paniculatum* extract one day before inoculation of ZYMV sap (T2), (iii) application of *C. paniculatum* extract on the same day as ZYMV applied (T3), (iv) application of *C. paniculatum* extract one day after ZYMV (T4), and no application of *C. paniculatum* extract (T0) (control).



Figure 1. Inoculum source of ZYMV isolates Bali on zucchini plants




Figure 2. Disease symptoms of ZYMV isolates Bali on zucchini plants: A. mosaic, B. yellowing with cupping, C. mosaic with cupping, and D. yellowing

The second factor was the concentration of inducer extract, which consisted three treatments, namely: (i) 10% of *C. paniculatum* extract (C1), (ii) 30% of *C. paniculatum* extract (C2), and (iii) 50% of *C. paniculatum* extract (C3). The treatments were combined to obtain (5 × 3) 15 combination treatments, and each combination treatment was repeated ten times to obtain 45 experimental units. Observation parameters included disease incidence, type of disease symptoms, disease severity, virus concentration, and loss of fruit yield. Data were analyzed using ANOVA according to a completely randomized design with a factorial pattern (5 × 3 × 10) with Duncan's follow-up test (DMRT) at 5% level.

Enzyme-linked immunosorbent assays (ELISA)

The detection of ZYMV virus was performed on the leaves of *C. pepo* plant four weeks after inoculation. Serological virus detection was carried out by enzyme-linked immunosorbent assay (DAS-ELISA)

using Potyvirus antiserum (DSMZ). Virus accumulation was quantitatively read using ELISA Reader model 550 (Bio-Rad, USA) at a wavelength of 405 nm.

RESULTS AND DISCUSSION

Incubation period

The incubation period of ZYMV in plants treated with *C. paniculatum* leaf extract was the most prolonged average incubation period (18.27 DAP) compared to the treatment without *C. paniculatum* leaf extract (C+) (3.60 DAP). Both factors (inoculation time and concentration of *C. paniculatum* leaf extract) exhibited a significantly longer incubation period when compared with the control treatment. The application of 50% *C. paniculatum* leaf extract one day before virus inoculation significantly slowed the incubation period of virus (21.00 DAP) compared with other treatments (Table 1). Kurnianingsih and Damayanti (2012) found that *C. paniculatum* leaf extract could slow the incubation period of BCMV in yardlong beans.

Symptoms of infection

ZYMV inoculation produced various symptoms, such as yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting (Figure 2 and Table 1). Plants treated with leaf extract one day after virus inoculation at concentrations of 10%, 30%, and 50% showed symptoms of yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting. The application of extract one day before inoculation of viral sap at concentrations of 30% and 50% only produced yellowing and yellowing with cupping symptoms.

Disease incidence

Plants treated with *C. paniculatum* leaf extract one day before sap virus inoculation showed the lowest (20%) disease incidence until the end of observation. The other treatments resulted in 40% - 60% disease prevalence. In control plants where no *C.*

paniculatum leaf extract was applied, disease incidence reached 100% (Table 4).

Disease severity

Clerodendrum paniculatum extract treatment significantly reduced the severity of disease. Among the treatments, the highest disease severity was recorded in extract application treatment one day after inoculation of viral smear. In comparison, the lowest was observed in the application of extract one day before the inoculation of virus (Table 2). The best time for application of *C. paniculatum* leaf extract was one day before ZYMV inoculation with a concentration of 50% because it could reduce disease severity up to 60.79% (Table 4). It was observed that *C. paniculatum* leaf extract suppressed the disease incidence and severity of ZYMV infection.

This is in accordance with Damayanti and Pebriyeni (2015), Kurnianingsih and Damayanti (2012), who reported that *C. paniculatum* leaf extract can suppress BCMV infection reaching 100% in yardlong beans.

Virus detection with ELISA

ELISA absorbance value (NAE) showed that no virus was detected in the samples given *C. paniculatum* leaf extract at a concentration of 30% and 50% one day before virus inoculation. NAE was not significantly different from the negative control (Table 4). Based on ELISA test, *C. pepo* showed presence (positive) of ZYMV in all treatments, except one day before virus sap inoculation with a concentration of 30 and 50%, *C. pepo* showed absence (negative) of ZYMV.

Table 1. Incubation period in ZYMV-infected zucchini plants in response to differences in application time and concentration of *Clerodendrum paniculatum* leaf extract

<i>C. paniculatum</i> leaf extract concentration (%)	Incubation period of ZYMV infection (DAP)				
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after virus sap inoculation
10	3.40 a	14.0 e	16.00 fg	6.20 b	6.00 b
30	3.60 a	15.2 f	17.80 h	8.60 d	7.20 c
50	3.80 a	16.8 g	21.00 i	9.40 d	8.40 d
Average	3.60 a	15.33 d	18.27 e	8.07 c	7.20 b

Note: day after planting (DAP); Different letters in the same row show a significant effect compared to the positive control (C+)

Table 2. Disease severity in ZYMV-infected zucchini plants in response to differences in application time and concentration of *Clerodendrum paniculatum* leaf extract

<i>C. paniculatum</i> leaf extract concentration (%)	Disease severity (%)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	66.37 j	39.96 e	31.80 bc	51.42 g	61.93 hij	50.30 c
30	64.30 ij	37.01 cd	27.33 ab	49.71 fg	58.78 hi	47.43 b
50	62.59 j	32.87 bc	23.59 a	45.14 ef	56.88 h	44.21 a
Average	64.42 e	36.61 b	27.57 a	48.76 c	59.19 d	

Note: Different letters in the same row show a significant effect compared to the positive control (C+)

Table 3. Flowering in zucchini plants in response to differences in application time and concentration of *Clerodendrum paniculatum* leaf extract

<i>C. paniculatum</i> leaf extract concentration (%)	Appearance of flower (DAP)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	37.00 f	33.60 bcd	32.80 b	35.80 e	36.80 f	35.20 b
30	37.20 f	32.60 b	33.80 bcd	33.20 bc	37.40 f	34.84 b
50	36.80 f	33.40 bc	30.80 a	34.60 cd	35.00 de	34.12 a
Average	37.00 c	33.20 a	32.47 a	34.53 b	36.40 c	

Note: day after planting (DAP); Different letters in the same row show a significant effect compared to the positive control (C+)

Table 4. Effect of application time and concentration of *Clerodendrum paniculatum* leaf extract on disease incidence, type of symptoms, suppression of ZYMV disease severity, and yield loss

Application time of <i>C. paniculatum</i> leaf extract	Extract concentration (%)	Disease incidence (%)	Type of disease symptom*	of Disease severity suppression (%)	ELISA results		Yield loss (%)
					Average of Reaction NAE**		
C+	10	100 (10/10)	M, Mc, St	-	0.418	+	58.42
	30	100 (10/10)	M, Mc, St	-	0.43	+	55.90
	50	60 (6/10)	Y, Yc, M	-	0.395	+	55.29
Extract application during seed soaking	10	60 (6/10)	M, Mc, St	39.77	0.557	+	35.56
	30	40 (4/10)	Y, Yc, Mc, St	42.58	0.486	+	33.69
	50	40 (4/10)	Y, Yc, Mc, St	47.70	0.458	+	37.41
Extract application one day before virus inoculation	10	40 (4/10)	Y, Yc	52.04	0.426	+	23.50
	30	20 (2/10)	Y, Yc	58.99	0.184	-	22.18
	50	20 (2/10)	Y, Yc	60.79	0.112	-	19.99
Extract application concurrently with viral inoculation	10	60 (6/10)	M, Mc, St	22.57	0.608	+	55.31
	30	40 (4/10)	M	23.25	0.515	+	40.74
	50	40 (4/10)	Y, Yc	26.96	0.489	+	43.93
Extract application one day after viral sap inoculation	10	60 (6/10)	Y, Yc, M, Mc, St	6.68	0.638	+	55.44
	30	60 (6/10)	M	8.96	0.588	+	55.19
	50	40 (4/10)	M	8.90	0.550	+	54.63

Note: C+ (without extract application). Type of disease symptom*: Y = yellowing, Yc = yellowing with cupping, M = mosaic, Mc = mosaic with cupping, St = stunting. Negative control: healthy plants (without virus inoculation and extract application). Positive control: sick plants (virus inoculation and without extract application). **NAE: Elisa Absorbance value

External



Ahmad Dwi Setyawan <smujo.id@gmail.com>

to me

Ni Putu Pandawani:

Thank you for submitting the manuscript, "High impact of pagoda flower leaves to suppress zucchini yield loss due to zucchini yellow mosaic virus infection" to Biodiversitas Journal of Biological Diversity. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Submission URL: <https://smujo.id/biodiv/authorDashboard/submission/11071>

Username: pandawani

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Ahmad Dwi Setyawan

Flowering and yield loss

Infection with ZYMV inhibits flowering in *C. pepo*. ZYMV-infected plants showed reduced flowering and yield loss in *C. pepo* (Kheder et al. 2017; Kone et al. 2017; Maghamnia et al. 2018; Maina et al. 2017). Plants infected with ZYMV first flowered at 37 DAP, while plants treated with *C. paniculatum* leaf extract one day before virus inoculation was flowered at 32 DAP (Table 3). Treatment of *C. paniculatum* leaf extract one day before virus inoculation on *C. pepo* resulted in a yield loss of 19.99% - 23.50% compared with other treatments. This indicated that applying *C. paniculatum* leaf extracts one day before ZYMV inoculation on *C. pepo* effectively reduced yield loss by 40%. Application of *C. paniculatum* leaf extract one day before virus inoculation increased the resistance of *C. pepo*. The low severity and virus accumulation in this treatment indicated the potential of the *C. paniculatum* leaf as an inducer of

systemic resistance of *C. pepo*. The mechanism of induction of systemic resistance of *C. pepo* by *C. paniculatum* leaf extract has been reported by Verma et al. (1998). The active compound in the form of 34 kDa protein in *C. paniculatum* flower leaf extract causes tobacco leaves to become immune to viruses. In addition, *C. paniculatum* flower leaf extract can suppress BCMV because it contains virus inhibitors and has antiviral activity (De Oliveira et al. 1993). The active compounds of the *C. paniculatum* flower are known as antiviral proteins and are known as ribosome-inactivating proteins (RIPs). RIPs are also present in the root and leaf extract of *M. jalapa* and are referred to as Mirabilis antiviral protein (MAP) (Verma et al. 1998). MAP can reach the ribosome's active site earlier than the virus to prevent viral infection at an early stage before the virus undergoes encapsidation (Vivanco et al. 1999).

In the present study *C. paniculatum* leaf extract showed 80% effectiveness in inhibiting ZYMV infection, whereas against BCMV and CMV, inhibition was observed to be 100% and 82.6%, respectively (Hersanti 2003; Kurnianingsih and Damayanti 2012). It indicates that the same plant extract showed different effectiveness against different viruses.

Application of *C. paniculatum* leaf extract one day before ZYMV inoculation with a concentration of 50% was able to reduce disease incidence and severity of, accelerate flowering, and reduce yield loss of *C. pepo* fruit compared to the control treatment. The results of this study enrich the information about the potential of *C. paniculatum* leaves as agent to control plant viruses.

ACKNOWLEDGEMENTS

Acknowledgments are conveyed to the farmer groups in Banjar Sandan Village, Baturiti, Tabanan, Bali for their assistance and support.

REFERENCES

- Aleem EEA, Rabie M, Fattouh FA. 2021. Molecular characterisation of zucchini yellow mosaic virus infecting *Cucurbita pepo* in Egypt. *Plant Prot Sci*. DOI: 10.17221/191/2020-PPS
- Bubici G, Navarro B, Carluccio AV, Ciuffo M, Serio FD, Fabrizio Cillo F. 2020. Genomic sequence variability of an Italian zucchini yellow mosaic virus isolate. *Eur J Plant Pathol* 156: 325-332.
- Coutts BA, Kehoe MA, Jones RAC. 2011. Minimising losses caused by zucchini yellow mosaic virus in vegetable cucurbit crops in tropical, subtropical and Mediterranean environments through cultural methods and host resistance. *Virus Res*. DOI: 10.1016/j.virusres.2011.04.015
- Coutts BA, Kehoe MAA, Jones RA. 2013. Zucchini yellow mosaic virus: Contact transmission, stability on surfaces, and inactivation with disinfectants. *Plant Dis*. DOI: 10.1094/PDIS-08-12-0769-RE
- Damayanti TA, Pebriyeni L. 2015. Barrier crop and pagoda leaf extract to control Bean common mosaic virus on yard long bean in the field. *J Hort*. 25: 238-245.
- De Oliveira MM, Sampaio MR, Noronha AB, Vicente M, Vianna S. 1993. Detection of antiviral and antitumoral fractions of *Chenopodium amaranticolor* leaf extract. *Microbios* 76: 213-221.
- Elbeshehy EF. 2017. Inhibitor activity of different medicinal plants extracts from *Thuja orientalis*, *Nigella sativa* L., *Azadirachta indica* and *Bougainvillea spectabilis* against Zucchini yellow mosaic virus (ZYMV) infecting *Citrullus lanatus*. *Biotechnol Equip*. DOI: 10.1080/13102818.2017.1279572
- Gal-On A. 2007. Zucchini yellow mosaic virus: Insect transmission and pathogenicity-The tails of two proteins. *Mol Plant Pathol*. DOI: 10.1111/j.1364-3703.2007.00381.x
- Hersanti. 2003. Testing the potential of extracts of 37 plant species as agents of inducing systemic resistance of red chili plants to Cucumber mosaic virus. *J Phytopathol Indones* 7: 54-58.
- Jaroszewska BH, Rymelska N, Borodynko N, Pospieszny H. 2013. Biological and molecular characterization of the Polish zucchini yellow mosaic virus isolates. *Acta Sci Pol Hortorum Cultus* 12: 75-85.
- Kheder AA, Taqwa M, Sulaiman TM, Ghanem GAM, Tohamy MR. 2017. Biological, serological and molecular characterization of Egyptian zucchini yellow mosaic virus isolate infecting squash plants in Fayoum governorate. *Egypt J Phytopathol*. DOI: 10.21608/ejp.2017.88570
- Kone N, Asare-Bediako E, Silue S, Kone D, Koita O, Wulf M, Stephan W. 2017. Influence of planting date on incidence and severity of viral disease on cucurbits under field condition. *Ann Agric Sci*. DOI: 10.1016/j.aos.2017.05.005

- Kurnianingsih L, Damayanti TA. 2012. Five plant extracts for suppressing Bean common mosaic virus infection on yard long bean. *J Phytopathol Indones*. DOI: 10.14692/jfi.8.6.155
- Lecoq H, Pitrat M, Cle'ment M. 1981. Identification et caracterisation d'un potyvirus provoquant la maladie du rabougrissement jaune du melon. *Agronomie* 1: 827-834.
- Lisa V, Boccardo G, D'Agostino G, Dellavalle G, D'Aquilio M. 1981. Characterization of a potyvirus that causes zucchini yellow mosaic. *Phytopathol* 71: 667-672.
- Maghamnia HR, Hajizadeh M, Azizi A. 2018. Complete genome sequence of zucchini yellow mosaic virus strain Kurdistan, Iran. *3 Biotech*. DOI: 10.1007/s13205-018-1177-3
- Maina S, Coutts BA, Edwards OR, Almeida L, Kehoe MA, Ximenes A, Jones RAC. 2017. Zucchini yellow mosaic virus populations from East Timorese and Northern Australian cucurbit crops: Molecular properties, genetic connectivity, and biosecurity implications. *Plant Dis*. DOI: 10.1094/PDIS-11-16-1672-RE
- Massumi H, Shaabani M, Heydarnejad J, Hosseini Pour AH, Rahimian H. 2011. Host range and phylogenetic analysis of Iranian isolates of zucchini yellow mosaic virus. *J Plant Pathol* 93: 187-193.
- Moradi Z, Mehrvar M, Nazifi E. 2019. Population genetic analysis of zucchini yellow mosaic virus based on the CI gene sequence. *J Cell Mol Res*. DOI: 10.22067/jcmr.v10i2.76133
- Nasr-Eldin MA, Hayam S, Abdelkader HS, Abo-Senna AS, Othman BA. 2016. Characterization and phylogenetic analysis of zucchini yellow mosaic virus infecting *Cucurbita pepo* in Egypt. *Am J Sci*. DOI: 10.7537/marsjas12031613
- Pandawani NP, Widnyana IK. 2021. Identification of virus causes of mosaic diseases in zucchini plants in the Bali Island of Indonesia. *J Tekirdag Agric Fac*. DOI: 10.33462/jotaf.707645
- Prendeville HR, Ye X, Morris J, Pilson D. 2012. Virus infections in wild plants are both frequent and often unapparent. *Am J Bot*. DOI: 10.3732/ajb.1100509
- Romay G, Lecoq H, Geraud-Pouey F, Chirinos DT, Desbiez C. 2014. Current status of cucurbit viruses in Venezuela and characterization of Venezuelan isolates of zucchini yellow mosaic virus. *Plant Pathol*. DOI: 10.1111/ppa.12072
- Simmons HE, Dunham JP, Zinn KE, Munkvold GP, Holmes EC, Stephenson AG. 2013. Zucchini yellow mosaic virus (ZYMV, Potyvirus): Vertical transmission, seed infection and cryptic infections. *Virus Res*. DOI: 10.1016/j.virusres.2013.06.016
- Simmons HE, Holmes EC, Gildow FE, Bothe-Goralczyk MA, Stephenson AG. 2011. Experimental verification of seed transmission of zucchini yellow mosaic virus. *Plant Dis*. DOI: 10.1094/PDIS-11-10-0843
- Spadotti D, Wassano D, Rezende J, Camargo LEA, Inoue-Nagata AK. 2015. Biological and molecular characterization of Brazilian isolates of zucchini yellow mosaic virus. *Sci Agric*. DOI: 10.1590/0103-9016-2014-0197
- Tymchyshyn O, Shevchenko T, Shevchenko O, Budzanivska I. 2017. Phylogenetic analysis of seed-transmitted isolate of zucchini yellow mosaic virus. *Bulletin of Taras Schevchenko National University of Kyiv* 2: 47-50.
- Valli AA, Gallo A, Rodamilans B, López-Moya JJ, García JA. 2017. The HCPPro from the Potyviridae family: an enviable multitasking helper component that every virus would like to have. *Mol Plant Pathol*. DOI: 10.1111/mpp.12553
- Verma HN, Baranwal VK, Srisavasta S. 1998. Antiviral substances of plant origin. In: Hadidi A, Khetarpal RK, Koganezawa H, editor. *Plant Virus Disease Control*. St. Paul Minnesota (US): APS Pr. Page 154-162.

Vivanco JM, Querci M, Salazar LF. 1999. Antiviral and anti-viroid activity of MAP containing extracts from *Mirabilis jalapa* roots. Plant Dis. DOI: 10.1094/PDIS.1999.83.12.1116

Wang D, Li G. 2017. Biological and molecular characterization of zucchini yellow mosaic virus isolates infecting melon in Xinjiang, China. Can J Plant Pathol. DOI: 10.1080/07060661.2017.1285356

=====

High impact of *Clerodendrum paniculatum* leaf extract to suppress zucchini yellow mosaic virus infection in zucchini plants

NI PUTU PANDAWANI^{1*}, LISTIHANI LISTIHANI¹, I KETUT WIDNYANA¹, PUTU EKA PASMIDI ARIATI¹, DEWA GEDE WIRYANGGA SELANGGA²

¹Program of Agrotechnology, Faculty of Agriculture and Business, Universitas Mahasaraswati Denpasar. Jl. Kamboja No.11A, Dangin Puri Kangin, Kec. Denpasar Utara, Kota Denpasar, Bali 80233. Tel./fax. +62- 361-227019, *email: pandawaniputu@unmas.ac.id

²Program of Agrotechnology, Faculty of Agriculture, Universitas Warmadewa. Jl. Terompong No.24, Sumerta Kelod, Kec. Denpasar Tim., Kota Denpasar, Bali 80239

Manuscript received: 2 May 2022. Revision accepted: xxx May 2022.

Abstract. Pandawani NP, Listihani L, Widnyana IK, Ariati PEP, Selangga DGW. 2022. High impact of *Clerodendrum paniculatum* leaf extract to suppress zucchini yellow mosaic virus infection in zucchini plants. *Biodiversitas* 23: xxxx. Zucchini yellow mosaic virus (ZYMV) is an important virus in zucchini plants (*Cucurbita pepo* L.). ZYMV infection is prevalent on *C. pepo* in Bali and is extremely difficult to eradicate. This study aimed to determine the efficacy of pagoda leaf extract (*Clerodendrum paniculatum* L.) in reducing yield loss in *C. pepo*. A factorial randomized block design with two variables was employed in this study. The application time of the *C. paniculatum* extract was the first variable, and concentration was the second. Plant extract was applied during seed immersion, one day before and after inoculation, and on the same day of virus inoculation. The concentrations of *C. paniculatum* extract employed were 10%, 30%, and 50%. The crude extract was sprayed onto the *C. pepo* leaves. Plants treated with extract one day after virus inoculation at 10%, 30%, and 50% concentrations exhibited yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting. While application of extract one day before virus inoculation at 30% and 50% concentrations resulted in yellowing and yellowing with cupping signs. Plants infected with ZYMV first showed flowering at 37 DAP, whereas plants treated with *C. paniculatum* leaf extract one day before virus inoculation started flowering at 32 DAP. Application of *C. paniculatum* leaf extract one day before ZYMV inoculation on *C. pepo* effectively suppressed 40% yield loss. The findings of present investigation revealed information about the potential of *C. paniculatum* leaf extract as a plant virus control agent through the mechanism of induction in *C. pepo*.

Keywords: Flowering, pagoda leaf extract, yield loss, zucchini, ZYMV

Abbreviations: Zucchini yellow mosaic virus (ZYMV)

INTRODUCTION

ZYMV is one of the most critical viruses causing mosaic disease in Cucurbitaceae plants worldwide, which could cause yield reduction (Simmons et al. 2011; Simmons et al. 2013). The incidence of viral disease due to ZYMV infection in cucumber plants could reach 100%, with varying attack intensities (Tymchyshyn et al. 2017). The loss due to viral infection depends on the time of infection and can result in yield losses of up to 100% (Coutts et al. 2011). Plants become very susceptible to viral infection at young age, affecting the high incidence of disease. When young plants become infected, the incubation period of virus is shorter, and the virus distribution and translocation process is accelerated (Coutts et al. 2013). Host range is one way of determining the

biological nature of a virus (Spadotti et al. 2015). The most frequently employed host range test plants for ZYMV are *Gomphrena globosa* L., *Chenopodium amaranticolor* Coste et Reyn., *C. quinoa* L., *Citrullus lanatus*, *Cucumis melo* L., *Cucumis sativus* L., *Luffa acutangula*, *Momordica charantia*, *Sechium edule*, *Phaseolus vulgaris* L., *Vigna sinensis*, *Capsicum annuum*, *Datura stramonium*, *Lycopersicon esculentum*, *Nicotiana tabacum* L. cv. White barley, *N. benthamiana*, and *Physalis floridana* (Jaroszewska et al. 2013; Massumi et al. 2013; Spadotti et al. 2015).

Zucchini yellow mosaic virus (ZYMV) has a one-molecule positive senseRNA genome of approximately 10 kb and encodes a poly-protein that is proteolytically processed into mature protein; P1 (protease), HC (helper component/protease), P3, 6K1, CI (cylinder inclusion),

6K2, NIa (nuclear inclusion a), VPg (viral protein-associated genome), NIB (nuclear inclusion b) and CP (Gal-On 2007; Moradi et al. 2019). Until now, four out of ten potyviral proteins have been identified in ZYMV-infected plants, namely P1, HC-Pro, cylindrical inclusions, cylindrical inclusions (CI) pinwheel type 1, and CP (Valli et al. 2017; Maghamnia et al. 2018). Other proteins (Third protein (P3), 6K1, 6K2, viral genome-linked protein (VPg), Nuclear Inclusion A (NIa), and NIB replicas) have not been identified in ZYMV-infected plants and are based on potyvirus sequence homology.

ZYMV was first reported in Italy in 1973 (Lisa et al. 1981) and then France (Lecoq et al. 1981). ZYMV distribution has been found in Africa, America, Asia, Europe, the Middle East, and Oceania (Spadotti et al. 2015; Bubici et al. 2020). In Egypt, ZYMV has been associated with significant yield losses in many cucurbit crops, including cucumbers, melons, pumpkins, marrow vegetables, watermelons, and especially zucchini (*Cucurbita pepo* L.) (Aleem et al. 2021). ZYMV infection result in symptoms, such as mosaicism, yellowing, and eventually "shoeing" on the leaves. The fruit is stunted, twisted, and distorted, resulting in reduced yield and inability to market the product, especially chayote zucchini (Massumi et al. 2011). ZYMV is commonly transferred in two ways: horizontally via aphid vectors and vertically by transmission from the first generation of ZYMV-infected seeds to the following generation of ZYMV-infected seeds (Simmons et al. 2011; Wang and Li 2017). Several species of aphids are ZYMV vectors that transmit viruses non-persistently, such as *A. gossypii*, *Myzus persicae* (Maina et al. 2017; Romay et al. 2014), *Asyrthosiphon pisum*, *A. kondoi*, *Aphis craccivora*, *A. citricola*, *A. middletonii*, *A. spiraeicola*, *Macrosiphum euphorbiae*, *Toxoptera aurantii*, and *Uroleucon ambrosiae* (Spadotti et al. 2015). ZYMV can also be transmitted mechanically easily through cutting tools that have been contaminated with the virus (Nasr-Eldin et al. 2016; Prendeville et al. 2012).

Viral diseases can be controlled by eradicating insect vectors, quarantining, and planting healthy and virus-free seeds. Additionally, the usage of antiviral compounds derived from plant extracts has been proven to be effective against a variety of viruses due to their ribosome-inactivating proteins (RIPs) (Gal-On 2007). One of the agents capable of inducing systemic resistance in plants is a plant extract (Elbeshehy 2017). The effective control of viral diseases is the use of resistant varieties, but there are not many commercial virus-resistant cultivars available.

Systemic resistance of a plant can be activated by inducing resistance genes present in plants by utilizing resistance-inducing agents (Elbeshehy 2017). One of the agents that induce systemic resistance of plants is plant extracts (Gal-On 2007). Spinach leaf extract (*Amaranthus spinosus*), four o'clock flower leaf (*Mirabilis jalapa*), and pagoda leaf (*Clerodendrum paniculatum* L.) were reported to induce systemic resistance to anthracnose and CMV pathogens in chili (Hersanti 2003). Other plant extract such as *Chenopodium amaranticolor* is reported to have antiviral activity for the tobacco mosaic virus (TMV) and antitumor for Ehrlich tumor (EA) (De Oliveira et al. 1993).

Extracts of *C. paniculatum* leaf, thorn spinach, four o'clock flower, *C. amaranticolor*, and sambiloto are able to suppress BCMV infection in yardlong bean plants (Kurnianingsih and Damayanti 2012).

The utilization of efficacious plant extracts have been used to benefit human health, but are still of little use in plant protection. Therefore, it is necessary to explore potential plant species to control pathogens, including viruses. The objective of this study was to test the effectiveness of plant extract to suppress ZYMV.

MATERIALS AND METHODS

Preparation of zucchini plant (*Cucurbita pepo* L.)

The seeds of the Carisa cultivar *C. pepo* were sown on plastic composite trays containing sterile growth media of a mixture of soil and compost. After 15 days, the grown seedlings were transplanted into plastic pots measuring 35 cm × 30 cm containing 5 kg of sterile planting medium, i.e. soil and manure (2:1 ratio). Each pot was planted with three seeds, and after one week of growth, one healthy plant was selected for the research object. Plants were irrigated daily with appropriate volume.

Propagation of ZYMV inoculum

The ZYMV inoculum was obtained from the Laboratory of the Faculty of Agriculture, Mahasaraswati University in Denpasar, then confirmed its nucleic acid sequence via DNA sequencing and designated it as ZYMV-zucchini isolate Bali (Pandawani and Widnyana 2021) (Figure 1). The inoculum was propagated mechanically by inoculating ZYMV inoculum on *C. pepo* two weeks after planting. Mechanical transmission of ZYMV was carried out by grinding sick leaves (source of inoculum) in phosphate buffer pH 7 with a ratio of 1:10 (w/v). The sap was then applied to the leaves of healthy plants which were treated with 600 mesh carborundum. After inoculation, leaves were rinsed with running water. The plants were then reared in a greenhouse, and symptomatic young leaves were used as a source of inoculum in this study.

Preparation and application of *Clerodendrum paniculatum* extract

The leaves of *C. paniculatum* was obtained from the area around Baturiti, Tabanan, Bali, Indonesia. *C. paniculatum* extract was prepared through a drying process to obtain a simplicia form. The simplicia was then macerated to produce a concentrated leaf extract with a concentration of 100% (Verma et al. 1998). Application of *C. paniculatum* extract to *C. pepo* was carried out according to the application time determined with a volume of 100 mL per plant. The *C. paniculatum* extract was applied on *C. pepo* leaves after three weeks of transplanting.

Inoculation of ZYMV in zucchini plants (*Cucurbita pepo* L.)

Three-week-old *C. pepo* were mechanically inoculated with ZYMV. Virus inoculum was prepared by grinding

symptomatic young leaves of *C. pepo* plant (source of ZYMV inoculum) in 1:10 (w/v) phosphate buffer. The virus was inoculated by applying the inoculum on the first leaf of *C. pepo* plant, which was sprinkled with 600 mesh carborundum.

Experiment design and data analysis

This study was conducted in a completely randomized design with a factorial pattern. The first factor was the time of application of the inducer extract, which consisted of five treatments, namely: (i) application of *C. paniculatum* extract during seed soaking (T1), (ii) application time of *C. paniculatum* extract one day before inoculation of ZYMV sap (T2), (iii) application of *C. paniculatum* extract on the same day as ZYMV applied (T3), (iv) application of *C. paniculatum* extract one day after ZYMV (T4), and no application of *C. paniculatum* extract (T0) (control).



Figure 1. Inoculum source of ZYMV isolates Bali on zucchini plants




Figure 2. Disease symptoms of ZYMV isolates Bali on zucchini plants: A. mosaic, B. yellowing with cupping, C. mosaic with cupping, and D. yellowing

The second factor was the concentration of inducer extract, which consisted three treatments, namely: (i) 10% of *C. paniculatum* extract (C1), (ii) 30% of *C. paniculatum* extract (C2), and (iii) 50% of *C. paniculatum* extract (C3). The treatments were combined to obtain (5 × 3) 15 combination treatments, and each combination treatment was repeated ten times to obtain 45 experimental units. Observation parameters included disease incidence, type of disease symptoms, disease severity, virus concentration, and loss of fruit yield. Data were analyzed using ANOVA according to a completely randomized design with a factorial pattern (5 × 3 × 10) with Duncan's follow-up test (DMRT) at 5% level.

Enzyme-linked immunosorbent assays (ELISA)

The detection of ZYMV virus was performed on the leaves of *C. pepo* plant four weeks after inoculation. Serological virus detection was carried out by enzyme-linked immunosorbent assay (DAS-ELISA) using Potyvirus antiserum (DSMZ). Virus accumulation was quantitatively read using ELISA Reader model 550 (Bio-Rad, USA) at a wavelength of 405 nm.

RESULTS AND DISCUSSION

Incubation period

The incubation period of ZYMV in plants treated with *C. paniculatum* leaf extract was the most prolonged average incubation period (18.27 DAP) compared to the treatment without *C. paniculatum* leaf extract (C+) (3.60 DAP). Both factors (inoculation time and concentration of

C. paniculatum leaf extract) exhibited a significantly longer incubation period when compared with the control treatment. The application of 50% *C. paniculatum* leaf extract one day before virus inoculation significantly slowed the incubation period of virus (21.00 DAP) compared with other treatments (Table 1). Kurnianingsih and Damayanti (2012) found that *C. paniculatum* leaf extract could slow the incubation period of BCMV in yardlong beans.

Symptoms of infection

ZYMV inoculation produced various symptoms, such as yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting (Figure 2 and Table 1). Plants treated with leaf extract one day after virus inoculation at concentrations of 10%, 30%, and 50% showed symptoms of yellowing, yellowing with cupping, mosaic, mosaic with cupping, and stunting. The application of extract one day before inoculation of viral sap at concentrations of 30% and 50% only produced yellowing and yellowing with cupping symptoms.

Disease incidence

Plants treated with *C. paniculatum* leaf extract one day before sap virus inoculation showed the lowest (20%) disease incidence until the end of observation. The other treatments resulted in 40% - 60% disease prevalence. In control plants where no *C. paniculatum* leaf extract was applied, disease incidence reached 100% (Table 4).

Disease severity

Clerodendrum paniculatum extract treatment significantly reduced the severity of disease. Among the treatments, the highest disease severity was recorded in extract application treatment one day after inoculation of viral smear. In comparison, the lowest was observed in the application of extract one day before the inoculation of virus (Table 2). The best time for application of *C. paniculatum* leaf extract was one day before ZYMV inoculation with a concentration of 50% because it could reduce disease severity up to 60.79% (Table 4). It was observed that *C. paniculatum* leaf extract suppressed the disease incidence and severity of ZYMV infection. This is in accordance with Damayanti and Pebriyeni (2015), Kurnianingsih and Damayanti (2012), who reported that *C. paniculatum* leaf extract can suppress BCMV infection reaching 100% in yardlong beans.

Virus detection with ELISA

ELISA absorbance value (NAE) showed that no virus was detected in the samples given *C. paniculatum* leaf extract at a concentration of 30% and 50% one day before virus inoculation. NAE was not significantly different from the negative control (Table 4). Based on ELISA test, *C. pepo* showed presence (positive) of ZYMV in all treatments, except one day before virus sap inoculation with a concentration of 30 and 50%, *C. pepo* showed absence (negative) of ZYMV.

Table 1. Incubation period in ZYMV-infected zucchini plants in response to differences in application time and concentration of *Clerodendrum paniculatum* leaf extract

<i>C. paniculatum</i> leaf extract concentration (%)	Incubation period of ZYMV infection (DAP)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	3.40 a	14.0 e	16.00 fg	6.20 b	6.00 b	9.12 a
30	3.60 a	15.2 f	17.80 h	8.60 d	7.20 c	10.48 b
50	3.80 a	16.8 g	21.00 i	9.40 d	8.40 d	11.88 c
Average	3.60 a	15.33 d	18.27 e	8.07 c	7.20 b	

Note: day after planting (DAP); Different letters in the same row show a significant effect compared to the positive control (C+)

Table 2. Disease severity in ZYMV-infected zucchini plants in response to differences in application time and concentration of *Clerodendrum paniculatum* leaf extract

<i>C. paniculatum</i> leaf extract concentration (%)	Disease severity (%)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	66.37 j	39.96 e	31.80 bc	51.42 g	61.93 hij	50.30 c
30	64.30 ij	37.01 cd	27.33 ab	49.71 fg	58.78 hi	47.43 b
50	62.59 j	32.87 bc	23.59 a	45.14 ef	56.88 h	44.21 a
Average	64.42 e	36.61 b	27.57 a	48.76 c	59.19 d	

Note: Different letters in the same row show a significant effect compared to the positive control (C+)

Table 3. Flowering in zucchini plants in response to differences in application time and concentration of *Clerodendrum paniculatum* leaf extract

<i>C. paniculatum</i> leaf extract concentration (%)	Appearance of flower (DAP)					Average effect of extract concentration
	No extract application (Positive control, C+)	Extract application during seed soaking	Extract application one day before virus sap inoculation	Extract application concurrently with viral sap inoculation	Extract application one day after viral sap inoculation	
10	37.00 f	33.60 bcd	32.80 b	35.80 e	36.80 f	35.20 b
30	37.20 f	32.60 b	33.80 bcd	33.20 bc	37.40 f	34.84 b
50	36.80 f	33.40 bc	30.80 a	34.60 cd	35.00 de	34.12 a
Average	37.00 c	33.20 a	32.47 a	34.53 b	36.40 c	

Note: day after planting (DAP); Different letters in the same row show a significant effect compared to the positive control (C+)

Table 4. Effect of application time and concentration of *Clerodendrum paniculatum* leaf extract on disease incidence, type of symptoms, suppression of ZYMV disease severity, and yield loss

Application time of <i>C. paniculatum</i> leaf extract	Extract concentration (%)	Disease incidence (%)	Type of disease symptom*	Disease severity suppression (%)	ELISA results		Yield loss (%)
					Average of NAE**	Reaction	
C+	10	100 (10/10)	M, Mc, St	-	0.418	+	58.42
	30	100 (10/10)	M, Mc, St	-	0.43	+	55.90
	50	60 (6/10)	Y, Yc, M	-	0.395	+	55.29
Extract application during seed soaking	10	60 (6/10)	M, Mc, St	39.77	0.557	+	35.56
	30	40 (4/10)	Y, Yc, Mc, St	42.58	0.486	+	33.69
	50	40 (4/10)	Y, Yc Mc, St	47.70	0.458	+	37.41
Extract application one day before virus sap inoculation	10	40 (4/10)	Y, Yc	52.04	0.426	+	23.50
	30	20 (2/10)	Y, Yc	58.99	0.184	-	22.18
	50	20 (2/10)	Y, Yc	60.79	0.112	-	19.99
Extract application concurrently with viral sap inoculation	10	60 (6/10)	M, Mc, St	22.57	0.608	+	55.31
	30	40 (4/10)	M	23.25	0.515	+	40.74
	50	40 (4/10)	Y, Yc	26.96	0.489	+	43.93
Extract application one day after viral sap inoculation	10	60 (6/10)	Y, Yc, M, Mc, St	6.68	0.638	+	55.44
	30	60 (6/10)	M	8.96	0.588	+	55.19
	50	40 (4/10)	M	8.90	0.550	+	54.63

Note: C+ (without extract application). Type of disease symptom*: Y = yellowing, Yc = yellowing with cupping, M = mosaic, Mc = mosaic with cupping, St = stunting. Negative control: healthy plants (without virus inoculation and extract application). Positive control: sick plants (virus inoculation and without extract application). **NAE: Elisa Absorbance value

Flowering and yield loss

Infection with ZYMV inhibits flowering in *C. pepo*. ZYMV-infected plants showed reduced flowering and yield loss in *C. pepo* (Kheder et al. 2017; Kone et al. 2017; Maghamnia et al. 2018; Maina et al. 2017). Plants infected with ZYMV first flowered at 37 DAP, while plants treated with *C. paniculatum* leaf extract one day before virus inoculation was flowered at 32 DAP (Table 3). Treatment of *C. paniculatum* leaf extract one day before virus inoculation on *C. pepo* resulted in a yield loss of 19.99% - 23.50% compared with other treatments. This indicated that applying *C. paniculatum* leaf extracts one day before ZYMV inoculation on *C. pepo* effectively reduced yield loss by 40%. Application of *C. paniculatum* leaf extract one day before virus inoculation increased the resistance of *C. pepo*. The low severity and virus accumulation in this treatment indicated the potential of the *C. paniculatum* leaf as an inducer of systemic resistance of *C. pepo*. The mechanism of induction of systemic resistance of *C. pepo* by *C. paniculatum* leaf extract has been reported by Verma

et al. (1998). The active compound in the form of 34 kDa protein in *C. paniculatum* flower leaf extract causes tobacco leaves to become immune to viruses. In addition, *C. paniculatum* flower leaf extract can suppress BCMV because it contains virus inhibitors and has antiviral activity (De Oliveira et al. 1993). The active compounds of the *C. paniculatum* flower are known as antiviral proteins and are known as ribosome-inactivating proteins (RIPs). RIPs are also present in the root and leaf extract of *M. jalapa* and are referred to as Mirabilis antiviral protein (MAP) (Verma et al. 1998). MAP can reach the ribosome's active site earlier than the virus to prevent viral infection at an early stage before the virus undergoes encapsidation (Vivanco et al. 1999).

In the present study *C. paniculatum* leaf extract showed 80% effectiveness in inhibiting ZYMV infection, whereas against BCMV and CMV, inhibition was observed to be 100% and 82.6%, respectively (Hersanti 2003; Kurnianingsih and Damayanti 2012). It indicates that the

same plant extract showed different effectiveness against different viruses.

Application of *C. paniculatum* leaf extract one day before ZYMV inoculation with a concentration of 50% was able to reduce disease incidence and severity of, accelerate flowering, and reduce yield loss of *C. pepo* fruit compared to the control treatment. The results of this study enrich the information about the potential of *C. paniculatum* leaves as agent to control plant viruses.

ACKNOWLEDGEMENTS

Acknowledgments are conveyed to the farmer groups in Banjar Sandan Village, Baturiti, Tabanan, Bali for their assistance and support.

REFERENCES

- Aleem EEA, Rabie M, Fattouh FA. 2021. Molecular characterisation of zucchini yellow mosaic virus infecting *Cucurbita pepo* in Egypt. *Plant Prot Sci*. DOI: 10.17221/191/2020-PPS
- Bubici G, Navarro B, Carluccio AV, Ciuffo M, Serio FD, Fabrizio Cillo F. 2020. Genomic sequence variability of an Italian zucchini yellow mosaic virus isolate. *Eur J Plant Pathol* 156: 325-332.
- Coutts BA, Kehoe MA, Jones RAC. 2011. Minimising losses caused by zucchini yellow mosaic virus in vegetable cucurbit crops in tropical, subtropical and Mediterranean environments through cultural methods and host resistance. *Virus Res*. DOI: 10.1016/j.virusres.2011.04.015
- Coutts BA, Kehoe MAA, Jones RA. 2013. Zucchini yellow mosaic virus: Contact transmission, stability on surfaces, and inactivation with disinfectants. *Plant Dis*. DOI: 10.1094/PDIS-08-12-0769-RE
- Damayanti TA, Pebriyeni L. 2015. Barrier crop and pagoda leaf extract to control Bean common mosaic virus on yard long bean in the field. *J Hort*. 25: 238-245.
- De Oliveira MM, Sampaio MR, Noronha AB, Vicente M, Vianna S. 1993. Detection of antiviral and antitumoral fractions of *Chenopodium amaranticolor* leaf extract. *Microbios* 76: 213-221.
- Elbeshy EF. 2017. Inhibitor activity of different medicinal plants extracts from *Thuja orientalis*, *Nigella sativa* L., *Azadirachta indica* and *Bougainvillea spectabilis* against Zucchini yellow mosaic virus (ZYMV) infecting *Citrullus lanatus*. *Biotechnol Biotechnol Equip*. DOI: 10.1080/13102818.2017.1279572
- Gal-On A. 2007. Zucchini yellow mosaic virus: Insect transmission and pathogenicity-The tails of two proteins. *Mol Plant Pathol*. DOI: 10.1111/j.1364-3703.2007.00381.x
- Hersanti. 2003. Testing the potential of extracts of 37 plant species as agents of inducing systemic resistance of red chili plants to Cucumber mosaic virus. *J Phytopathol Indones* 7: 54-58.
- Jaroszewska BH, Rymelska N, Borodynyko N, Pospieszny H. 2013. Biological and molecular characterization of the Polish zucchini yellow mosaic virus isolates. *Acta Sci Pol Hortorum Cultus* 12: 75-85.
- Kheder AA, Taqwa M, Sulaiman TM, Ghanem GAM, Tohamy MR. 2017. Biological, serological and molecular characterization of Egyptian zucchini yellow mosaic virus isolate infecting squash plants in Fayoum governorate. *Egypt J Phytopathol*. DOI: 10.21608/ejp.2017.88570
- Kone N, Asare-Bediako E, Silue S, Kone D, Koita O, Wulf M, Stephan W. 2017. Influence of planting date on incidence and severity of viral disease on cucurbits under field condition. *Ann Agric Sci*. DOI: 10.1016/j.aosas.2017.05.005
- Kurnianingsih L, Damayanti TA. 2012. Five plant extracts for suppressing Bean common mosaic virus infection on yard long bean. *J Phytopathol Indones*. DOI: 10.14692/jfi.8.6.155
- Lecoq H, Pitrat M, Cle'ment M. 1981. Identification et caracterisation d'un potyvirus provoquant la maladie du rabougrissement jaune du melon. *Agronomie* 1: 827-834.
- Lisa V, Boccardo G, D'Agostino G, Dellavalle G, D'Aquilio M. 1981. Characterization of a potyvirus that causes zucchini yellow mosaic. *Phytopathol* 71: 667-672.
- Maghamnia HR, Hajizadeh M, Azizi A. 2018. Complete genome sequence of zucchini yellow mosaic virus strain Kurdistan, Iran. *3 Biotech*. DOI: 10.1007/s13205-018-1177-3
- Maina S, Coutts BA, Edwards OR, Almeida L, Kehoe MA, Ximenes A, Jones RAC. 2017. Zucchini yellow mosaic virus populations from EastTimorese and Northern Australian cucurbit crops: Molecular properties, genetic connectivity, and biosecurity implications. *Plant Dis*. DOI: 10.1094/PDIS-11-16-1672-RE
- Massumi H, Shaabani M, Heydarnejad J, Hosseini Pour AH, Rahimian H. 2011. Host range and phylogenetic analysis of Iranian isolates of zucchini yellow mosaic virus. *J Plant Pathol* 93: 187-193.
- Moradi Z, Mehrvar M, Nazifi E. 2019. Population genetic analysis of zucchini yellow mosaic virus based on the CI gene sequence. *J Cell Mol Res*. DOI: 10.22067/jcmr.v10i2.76133
- Nasr-Eldin MA, Hayam S, Abdelkader HS, Abo-Senna AS, Othman BA. 2016. Characterization and phylogenetic analysis of zucchini yellow mosaic virus infecting *Cucurbita pepo* in Egypt. *Am J Sci*. DOI: 10.7537/marsjas12031613
- Pandawani NP, Widnyana IK. 2021. Identification of virus causes of mosaic diseases in zucchini plants in the Bali Island of Indonesia. *J Tekirdag Agric Fac*. DOI: 10.33462/jotaf.707645
- Prendeville HR, Ye X, Morris J, Pilsion D. 2012. Virus infections in wild plants are both frequent and often unapparent. *Am J Bot*. DOI: 10.3732/ajb.1100509
- Romay G, Lecoq H, Geraud-Pouey F, Chirinos DT, Desbiez C. 2014. Current status of cucurbit viruses in Venezuela and characterization of Venezuelan isolates of zucchini yellow mosaic virus. *Plant Pathol*. DOI: 10.1111/ppa.12072
- Simmons HE, Dunham JP, Zinn KE, Munkvold GP, Holmes EC, Stephenson AG. 2013. Zucchini yellow mosaic virus (ZYMV, Potyvirus): Vertical transmission, seed infection and cryptic infections. *Virus Res*. DOI: 10.1016/j.virusres.2013.06.016
- Simmons HE, Holmes EC, Gildow FE, Bothe-Goralczyk MA, Stephenson AG. 2011. Experimental verification of seed transmission of zucchini yellow mosaic virus. *Plant Dis*. DOI: 10.1094/PDIS-11-10-0843
- Spadotti D, Wassano D, Rezende J, Camargo LEA, Inoue-Nagata AK. 2015. Biological and molecular characterization of Brazilian isolates of zucchini yellow mosaic virus. *Sci Agric*. DOI: 10.1590/0103-9016-2014-0197
- Tymchysyn O, Shevchenko T, Shevchenko O, Budzanivska I. 2017. Phylogenetic analysis of seed-transmitted isolate of zucchini yellow mosaic virus. *Bulletin of Taras Schevchenko National University of Kyiv* 2: 47-50.
- Valli AA, Gallo A, Rodamilans B, López-Moya JJ, García JA. 2017. The HCPro from the Potyviridae family: an enviable multitasking helper component that every virus would like to have. *Mol Plant Pathol*. DOI: 10.1111/mpp.12553
- Verma HN, Baranwal VK, Srisavasta S. 1998. Antiviral substances of plant origin. In: Hadidi A, Khetarpal RK, Koganezawa H, editor. *Plant Virus Disease Control*. St. Paul Minnesota (US): APS Pr. Page 154-162.
- Vivanco JM, Querci M, Salazar LF. 1999. Antiviral and antiviroid activity of MAP containing extracts from *Mirabilis jalapa* roots. *Plant Dis*. DOI: 10.1094/PDIS.1999.83.12.1116
- Wang D, Li G. 2017. Biological and molecular characterization of zucchini yellow mosaic virus isolates infecting melon in Xinjiang, China. *Can J Plant Pathol*. DOI: 10.1080/07060661.2017.1285356

Dear Biodiversitas team, Thank you for your information. I have corrected the article according to suggestions from reviewers. can this article be included in the full article? not put into short communication. Best regards, Ni putu pandawani [pandawani, A-11097-Research Results-1052925-1-4-20220520_R2.doc](#)

pandawani
2022-05-25
11:21 AM

BILLING

dewinurpratiwi
2022-05-25 04:50 AM

BILLING

[Close Panel](#)

Participants

- DEWI NUR PRATIWI (dewinurpratiwi)
- Ni Putu Pandawani (pandawani)

Messages

Note

Dear Author(s),

From

dewinurpratiwi

Note

From

2022-05-25 04:50 AM

Kindly find attached an invoice for the publication of your manuscript.

[dewinurpratiwi, 3978.NI PUTU PANDAWANI.pdf](#)

INVOICE FOR AUTHOR

=====

[biodiv] Editor Decision

External

Inbox

Anisa Septiasari

Mon, May 30, 12:52 PM (22 hours ago)

to me, LISTIHANI

NI PUTU PANDAWANI, LISTIHANI LISTIHANI, I KETUT WIDNYANA, PUTU EKA PASMIDI
ARIATI, DEWA GEDE WIRYANGGA SELANGGA:

We have reached a decision regarding your submission to Biodiversitas Journal of
Biological Diversity, "High impact of Clerodendrum paniculatum leaf extract to suppress
zucchini yellow mosaic virus infection in zucchini plants".

Our decision is to: Accept Submission

30 Mei 2022 -Biodiv jurnal

My Assigned

Search

[New Submission](#)

Filters

Overdue

Incomplete

Stages

Submission

Review

Copyediting

Production

Sections

Articles

Activity

[Days since last activity](#)

Days since last activity

ID11097

PANDAWANI et al.

High impact of Clerodendrum paniculatum leaf extract to suppress zucchini yellow mosaic virus infection in zucchini plants

CopyeditingCurrently in the Copyediting stage.

1

Hide expanded details about PANDAWANI et al.

1Copyedited files submitted

0Open discussions

Last activity recorded on Monday, May 30, 2022.

11097 / **PANDAWANI et al.** / High impact of Clerodendrum paniculatum leaf extract to suppress zucchini yellow mosaic virus infection in zucchini plants

Library

WorkflowPublication

- [Submission](#)

- [Review](#)

- [Copyediting](#)

- [Production](#)

Copyediting Discussions

- [Add discussion](#)

Name	From	Last Reply	Replies	Closed
------	------	------------	---------	--------

No Items

Copyedited

- **Search**

Name	Date	Component
1053553-1 aseptiasari, D230618-Flowering - Pandawani=REV-check.doc	May 30, 2022	Article Text